

Beyond genotype: serotonin transporter epigenetic modification predicts human brain function

Yuliya S Nikolova¹, Karestan C Koenen², Sandro Galea², Chiou-Miin Wang³, Marianne L Seney⁴, Etienne Sibille⁴, Douglas E Williamson⁵ & Ahmad R Hariri¹

We examined epigenetic regulation in regards to behaviorally and clinically relevant human brain function. Specifically, we found that increased promoter methylation of the serotonin transporter gene predicted increased threat-related amygdala reactivity and decreased mRNA expression in postmortem amygdala tissue. These patterns were independent of functional genetic variation in the same region. Furthermore, the association with amygdala reactivity was replicated in a second cohort and was robust to both sampling methods and age.

The systematic integration of human molecular genetics and *in vivo* neuroimaging have contributed to our increasing understanding of how DNA sequence-based genetic variation shapes individual differences in brain function, complex behavioral traits and related risk for psychopathology¹. Parallel research in animal models has highlighted a critical role for non-sequence-based epigenetic variation in the emergence of individual differences in brain function and risk-related behavior². The importance of similar epigenetic mechanisms for behaviorally and clinically relevant brain function in humans has yet to be fully explored.

We used bisulfite sequencing to determine the percentage of methylation of the proximal promoter region of the serotonin transporter gene (*SLC6A4*) in saliva-derived DNA from a discovery cohort of 80 young adults and blood-derived DNA from an independent replication cohort of 96 adolescents. We targeted *SLC6A4* because it encodes the serotonin transporter, which modulates brain function and behavior by regulating the duration and intensity of synaptic serotonin signaling. Dysfunction of the serotonin transporter has also been implicated in the pathophysiology of mood and anxiety disorders³, and pharmacologic blockade of this molecule is the primary mode of treating these same disorders.

We focused our analyses on the 20 CpG sites closest to the transcription start site (TSS) of *SLC6A4* exhibiting substantial variability across individuals (Table 1 and Supplementary Fig. 1a,b). Additional proximal promoter sites were excluded, as we found

virtually no variability across individuals (Supplementary Table 1). In light of recent work suggesting that methylation immediately downstream of the TSS may also affect transcription⁴, we sampled additional CpG sites up to 119 base pairs downstream of the TSS, spanning exon 1 and intron 1 (Supplementary Table 1), whose effects were investigated in separate control analyses.

We evaluated the relationship between *SLC6A4* proximal promoter methylation and threat-related amygdala reactivity assayed using blood oxygen level-dependent (BOLD) functional magnetic resonance imaging (fMRI). We selected this neural phenotype as a measure of behaviorally and clinically relevant brain function because it is clearly involved in the emergence of both normal and pathologic emotional behaviors^{5,6}. Notably, these behaviors include responsiveness to environmental and social stress, which is associated not only with epigenetic modification⁷, but also with variability in serotonin signaling⁸. Moreover, there is now ample evidence linking variability in serotonin signaling with individual differences in amygdala reactivity^{5,9,10}.

In our discovery cohort, the percentage of methylation of the *SLC6A4* proximal promoter was positively correlated with threat-related amygdala reactivity in the left hemisphere (adjusted $R^2 = 0.067$, $b = 0.282$, $P = 0.011$; Fig. 1a,b). This effect was observed at a trend level in the right hemisphere (adjusted $R^2 = 0.032$, $b = 0.211$, $P = 0.060$). The percentage of methylation continued to account for significant variability in left amygdala reactivity even when controlling for possible effects of gender, age, early and recent life stress, and current psychiatric disorder (left hemisphere: $\Delta R^2 = 0.084$, $b = 0.292$, $P = 0.009$; right hemisphere: $\Delta R^2 = 0.045$, $b = 0.214$, $P = 0.060$). Similar results were obtained when using the top principal component (PC) capturing 24% of the methylation variance in the same region (Supplementary Tables 2 and 3, and Supplementary Fig. 2a,b). In an exploratory follow-up analysis, we probed the effects of individual CpG site methylation levels on these same phenotypes and found that CpG 14 (188 base pairs upstream of TSS) showed the strongest association effects across both hemispheres (Table 1).

Given prior work establishing predictive links between genetic variation and amygdala reactivity¹, we next compared the effect of *SLC6A4* proximal promoter methylation on amygdala reactivity to that of the serotonin transporter-linked polymorphic region (5-HTTLPR) and rs25531, which together define a functional tri-allelic polymorphism previously associated with variability in amygdala reactivity and responsiveness to stress³. *SLC6A4* methylation continued to predict amygdala reactivity even when the 5-HTTLPR/rs25531 genotype was accounted for alongside all other covariates (left hemisphere: $\Delta R^2 = 0.086$, $b = 0.296$, $P = 0.009$; right hemisphere: $\Delta R^2 = 0.043$, $b = 0.211$, $P = 0.066$). This suggests that

¹Laboratory of NeuroGenetics, Department of Psychology & Neuroscience, and Institute for Genome Sciences & Policy, Duke University, Durham, North Carolina, USA. ²Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, New York, USA. ³Department of Molecular Medicine, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA. ⁴Department of Psychiatry, Center for Neuroscience, University of Pittsburgh, Pittsburgh, Pennsylvania, USA. ⁵Department of Psychiatry, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA. Correspondence should be addressed to Y.S.N. (yuliya.nikolova@duke.edu).

Received 1 February; accepted 9 July; published online 3 August 2014; doi:10.1038/nn.3778

Table 1 Summary of results from linear regression models predicting *in vivo* amygdala reactivity and amygdala tissue *SLC6A4* mRNA from percentage methylation at each of the 20 individual proximal promoter CpG sites sampled across the discovery, replication and postmortem cohorts

CpG site	Distance to TSS	Discovery cohort (DNS)				Replication cohort (TAOS)				Postmortem	
		Left amygdala		Right amygdala		Left amygdala		Right amygdala		Amygdala <i>SLC6A4</i> mRNA	
		<i>b</i>	<i>p</i>	<i>b</i>	<i>p</i>	<i>b</i>	<i>p</i>	<i>b</i>	<i>p</i>	<i>b</i>	<i>p</i>
1	-69	0.163	0.149	0.180	0.11	0.112	0.278	-0.061	0.556	-0.073	0.707
2	-72	0.263	0.018	0.125	0.269	0.207	0.043	0.068	0.511	-0.043	0.837
3	-99	0.094	0.409	0.052	0.647	0.119	0.248	0.148	0.15	0.118	0.540
4	-112	0.224	0.046	0.279	0.012	0.107	0.297	0.130	0.208	-0.079	0.752
5	-133	0.102	0.368	0.138	0.223	0.229	0.025	0.070	0.500	-0.128	0.601
6	-135	0.224	0.046	0.260	0.02	0.129	0.209	-0.106	0.306	0.104	0.630
7	-139	0.069	0.544	0.191	0.089	0.164	0.109	-0.019	0.856	-0.271	0.169
8	-141	0.026	0.818	0.048	0.672	0.010	0.921	0.020	0.846	-0.006	0.975
9	-147	0.128	0.258	0.058	0.608	0.196	0.056	-0.056	0.587	0.022	0.920
10	-149	0.280	0.012*	0.225	0.045	0.134	0.193	0.085	0.411	0.070	0.708
11	-155	0.186	0.099	0.240	0.032	0.214	0.037	0.133	0.195	0.088	0.683
12	-170	0.162	0.151	0.144	0.204	0.264	0.009	0.070	0.496	-0.223	0.301
13	-174	0.245	0.028	0.219	0.051	0.273	0.007	0.213	0.037	0.071	0.702
14	-188	-0.029	0.799	0.287	0.010*	0.305	0.003*	0.219	0.032*	-0.378	0.039*
15	-190	0.157	0.164	0.164	0.146	0.218	0.033	0.097	0.345	-0.013	0.943
16	-195	0.094	0.407	-0.103	0.363	0.226	0.027	0.037	0.722	0.177	0.407
17	-200	0.006	0.955	-0.107	0.343	0.082	0.424	0.004	0.965	0.166	0.495
18	-207	0.148	0.189	0.091	0.423	0.226	0.027	0.014	0.893	-0.032	0.865
19	-209	0.119	0.294	-0.059	0.602	0.235	0.021	0.029	0.782	-0.082	0.671
20	-213	0.100	0.376	-0.057	0.616	0.115	0.263	-0.116	0.261	-0.729	0.472

Results from the *in vivo* imaging cohorts are not adjusted for covariates. In light of gender, age, pH, postmortem interval and RNA ratio effects in the postmortem cohort, the results for the postmortem findings are adjusted for covariates. The CpG site numbering scheme reflects the ordering of CpG site in this proximal promoter region and has no relation to any unique CpG site numerical identifiers. All significant *P* values are shown in bold.

*CpG site with strongest association for each phenotype.

there is a preponderance of epigenetic variation over sequence-based variation in regulatory regions of the same gene.

As a negative control, we examined the correlation between amygdala reactivity and methylation in other regions of *SLC6A4* and the *COMT* gene, which codes for an enzyme responsible for regulating catecholamine, but not serotonin, signaling. As expected, there were no significant correlations between left or right amygdala reactivity and percent methylation of either intron 1 or exon 1 of *SLC6A4* ($P > 0.10$). Similarly, there were also no significant correlations between reactivity and percent promoter methylation of *COMT* ($P > 0.50$; **Supplementary Table 4**). Thus, these data reveal a specific effect of *SLC6A4* proximal promoter methylation on amygdala reactivity. Given the specificity of these findings, we followed up the effects of these same 20 CpG sites in our replication cohort.


Consistent with the findings from our discovery cohort, we found that the percentage of promoter methylation of *SLC6A4*, this time assayed in DNA derived from peripheral blood, was positively correlated with left amygdala reactivity in our replication cohort (adjusted $R^2 = 0.104$, $b = 0.336$, $P = 0.001$; **Fig. 1c,d**). In fact, this effect of percent methylation was larger than that accounted for in our discovery cohort (10.4 versus 6.7%). The effect was again weaker in the right hemisphere (adjusted $R^2 = 0.007$, $b = 0.084$, $P = 0.42$). Such hemispheric

asymmetries are not uncommon in the imaging genetics literature^{9,10} and may reflect task-specific characteristics¹¹ or intrinsic differences in monoamine signaling between the two hemispheres¹².

Notably, the effects in our replication cohort remained unchanged when controlling for age, gender, early life stress, risk for psychiatric disorder, and 5-HTTLPR/rs25531 genotype (left hemisphere: $\Delta R^2 = 0.090$, $b = 0.312$, $P = 0.005$; right hemisphere: $\Delta R^2 = 0.002$, $b = 0.049$, $P = 0.662$). These results were confirmed using the first PC capturing 30.41% of all methylation variance in the region (**Supplementary Tables 5 and 6**, and **Supplementary Fig. 2c,d**). Finally, as in the discovery cohort, the single site showing strongest correlations with amygdala reactivity across hemispheres was CpG 14 (**Table 1**).

The direction of the effects that we observed *in vivo* (higher reactivity with greater percent methylation) is markedly consistent with that of prior imaging genetics^{5,9}, multi-modal PET/fMRI^{10,13} and

Figure 1 Effects of *SLC6A4* promoter methylation on amygdala reactivity. (a,c) Statistical parametric map illustrating mean bilateral threat-related amygdala reactivity across all participants in the discovery cohort (left: $x = -24$, $y = -8$, $z = -16$, $t = 10.29$, $P = 2.0095 \times 10^{-14}$, $k_E = 180$; right: $x = 30$, $y = -4$, $z = -20$, $t = 11.13$, $P < 0.00001$, $k_E = 203$; a) and replication cohort (left: $x = -20$, $y = -4$, $z = -18$, $t = 11.29$, $P < 0.00001$, $k_E = 197$; right: $x = 20$, $y = -4$, $z = -16$, $t = 11.80$, $P = 1.29 \times 10^{-13}$, $k_E = 205$; c). Activation clusters are overlaid onto canonical structural brain images in the coronal plane ($Y = -2$). (b,d) The average percent *SLC6A4* proximal promoter methylation was positively correlated with reactivity of the left amygdala in both the discovery (b) and replication cohorts (d). a.u. = arbitrary units. $P < 0.05$.






Figure 2 Effects of *SLC6A4* promoter methylation on gene expression. There was a negative correlation between percent methylation at CpG 14 and *SLC6A4* mRNA levels in amygdala tissue. Units on the y axis units are expressed as arbitrary signal ($2^{-\text{dCT}} \times 10,000$), square root transformed to normalize the distribution of the variable.

pharmacologic fMRI^{14,15} studies linking relatively increased serotonin signaling with increased amygdala reactivity, as well as with observations that methylation in or near promoter regions generally inhibits gene transcription⁴. In light of these data, and to gain further mechanistic insight into our *in vivo* findings, we examined the effect of percent methylation of the same 20 CpG sites sampled in our imaging cohorts on serotonin transporter mRNA levels in postmortem amygdala tissue from a third independent cohort of 34 individuals.

As expected, clinical and biochemical parameters (that is, diagnostic status, pH and RNA ratio) influenced mRNA levels ($P < 0.091$). Thus, these were included as covariates alongside age, gender and postmortem interval in all analyses involving postmortem data. When controlling for the effects of these parameters, there was no significant association between overall percent methylation and mRNA in amygdala tissue ($P = 0.699$). However, a site-specific investigation revealed a significant negative correlation ($P = 0.039$) between mRNA levels and percent methylation at CpG 14, which exhibited the strongest correlation with amygdala reactivity in both our imaging cohorts (Table 1 and Fig. 2). As with our *in vivo* imaging data, this epigenetic effect was further independent of 5-HTTLPR/rs25531 genotype ($P = 0.031$).

Collectively, our results suggest that methylation of the proximal promoter of human *SLC6A4* predicts threat-related amygdala reactivity, possibly reflecting decreased serotonin transporter gene expression and, consequently, reduced regional serotonin reuptake. Moreover, these epigenetic effects are independent of, and greater than, the effects of the 5-HTTLPR/rs25531 functional polymorphism near the same genomic region. Further demonstrating the independence of these genetic and epigenetic effects, none of the 20 proximal promoter CpG sites surveyed across the three cohorts overlapped the 5-HTTLPR, which is located 1,400 base pairs upstream of the TSS and primarily affects the distal promoter. In addition, the 5-HTTLPR/rs25531 genotype had no effect on proximal promoter methylation in any of the cohorts ($P > 0.30$).

Although we did not directly map methylation in peripheral tissues onto methylation levels in brain in the same cohort, cross-cohort convergence among tissues (saliva, blood, brain) is consistent with recent work demonstrating a substantial correlation between the blood and brain methylomes¹⁶. Furthermore, prior work has mapped variability in a similar range of *SLC6A4* promoter methylation values onto individual differences in serotonin biosynthesis *in vivo*¹⁷. Finally, the cross-tissue convergence that we observed suggests that two distinct types of readily assayed peripheral tissues (blood and saliva) could potentially be used as equally valid proxies of neural tissue. A notable limitation of our work is that, as a result of practical constraints, the postmortem tissue analysis was limited to the amygdala, although more informative differences in *SLC6A4* transcript levels are likely to

emerge in the dorsal raphe nucleus, where the serotonin transporter is more densely expressed. This limitation notwithstanding, our results demonstrate that meaningful links between the human epigenome and brain can be mapped using DNA derived from readily assayed peripheral tissues. In addition to encouraging careful consideration of the effects of promoter methylation in *SLC6A4* on behaviorally and clinically relevant brain function, we hope that our results will advance broader research on epigenetic mechanisms in the emergence of individual differences in human behavior and related risk for psychopathology.

METHODS

Methods and any associated references are available in the [online version of the paper](#).

Note: Any Supplementary Information and Source Data files are available in the online version of the paper.

ACKNOWLEDGMENTS

We thank B. Brigidi, K. Faig, A. Gorke, S. Jacobson, A. Knodt, B. Williams and K. Sugden for their assistance in DNS data collection and analysis, and J. Hanson for his assistance in figure preparation. The Duke Neurogenetics Study is supported by Duke University and National Institute on Drug Abuse grant DA03369. The Teen Alcohol Outcomes Study was supported by AA016274 and ongoing support from the Dielmann Family (D.E.W.). Y.S.N. receives support through a Howard Hughes Medical Institute International Student Research fellowship. A.R.H. receives support through National Institute on Drug Abuse grants DA033369 and DA031579. K.C.K. receives support through National Institute of Mental Health grants MH078928 and MH093612. E.S. receives support through National Institute of Mental Health grants MH084060 and MH077159.

AUTHOR CONTRIBUTIONS

Y.S.N. designed the study, participated in the collection of the neuroimaging and genetic data for the discovery cohort, conducted all of the statistical analyses, and wrote the manuscript with A.R.H. K.C.K. and S.G. designed and coordinated the methylation analyses in the discovery cohort. C.-M.W. developed and performed the methylation assays in the replication and postmortem cohorts. M.L.S. conducted the quantitative PCR in the postmortem cohort. E.S. designed the parent protocol, supervised quantitative PCR experiments and coordinated methylation analyses in the postmortem cohort. D.E.W. designed the parent protocol for the replication cohort and coordinated the methylation assays in both the replication and postmortem cohorts. A.R.H. designed the study, coordinated all analyses and wrote the manuscript with Y.S.N. He also designed the parent protocol for the discovery cohort and the neuroimaging protocol for the replication cohort. All of the authors provided feedback on the manuscript and approved its final version.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

Reprints and permissions information is available online at <http://www.nature.com/reprints/index.html>.

1. Hariri, A.R. *Annu. Rev. Neurosci.* **32**, 225–247 (2009).
2. Meaney, M.J. & Szyf, M. *Dialogues Clin. Neurosci.* **7**, 103–123 (2005).
3. Caspi, A., Hariri, A.R., Holmes, A., Uher, R. & Moffitt, T.E. *Am. J. Psychiatry* **167**, 509–527 (2010).
4. Brenet, F. *et al. PLoS ONE* **6**, e14524 (2011).
5. Fakra, E. *et al. Arch. Gen. Psychiatry* **66**, 33–40 (2009).
6. Abercrombie, H.C. *et al. Neuroreport* **9**, 3301–3307 (1998).
7. Mehta, D. *et al. Proc. Natl. Acad. Sci. USA* **110**, 16044–16049 (2013).
8. Chaouloff, F. *J. Psychopharmacol.* **14**, 139–151 (2000).
9. Hariri, A.R. *et al. Science* **297**, 400–403 (2002).
10. Fisher, P.M. *et al. Nat. Neurosci.* **9**, 1362–1363 (2006).
11. Baas, D., Aleman, A. & Kahn, R.S. *Brain Res. Brain Res. Rev.* **45**, 96–103 (2004).
12. Young, E.J. & Williams, C.L. *Behav. Neurosci.* **124**, 633–644 (2010).
13. Rhodes, R.A. *et al. J. Neurosci.* **27**, 9233–9237 (2007).
14. Bigos, K.L. *et al. Neuropsychopharmacology* **33**, 3221–3225 (2008).
15. Di Simplicio, M., Norbury, R., Reinecke, A. & Harmer, C.J. *Psychol. Med.* **44**, 241–252 (2013).
16. Tylee, D.S., Kawaguchi, D.M. & Glatt, S.J. *Am. J. Med. B Neuropsychiatr. Genet.* **162B**, 595–603 (2013).
17. Wang, D. *et al. PLoS ONE* **7**, e39501 (2012).

ONLINE METHODS

Discovery cohort. The first 91 Caucasian participants (47 women, mean age = 19.66 ± 1.36 years) to complete the ongoing Duke Neurogenetics Study (DNS) were selected for inclusion in analyses involving DNA methylation assays. The DNS assesses a range of behavioral and biological traits among young adult, student volunteers. All participants provided informed consent in accord with Duke University guidelines, and were in good general health. Two participants' samples displayed incorrect sequence patterns for our *SLC6A4* promoter assays as a result of unknown mutations or equipment dispensation error and were excluded from this analysis. Nine additional participants were excluded because of task non-compliance or response box failure, leaving a final sample of 80 individuals (42 women, mean age = 19.74 ± 1.33 years).

All participants were free of the following study exclusions: (1) medical diagnoses of cancer, stroke, diabetes requiring insulin treatment, chronic kidney or liver disease, or lifetime history of psychotic symptoms, (2) use of psychotropic, glucocorticoid or hypolipidemic medication, and (3) conditions affecting cerebral blood flow and metabolism (for example, hypertension). Diagnosis of any current DSM-IV Axis I disorder or select Axis II disorders (Antisocial Personality Disorder and Borderline Personality Disorder), assessed with the electronic Mini International Neuropsychiatric Interview¹⁸ and Structured Clinical Interview for the DSM-IV (SCID) substests¹⁹, respectively, was not an exclusion as the DNS seeks to establish broad variability in multiple behavioral phenotypes related to psychopathology. However, all participants were medication-free at the time of the study. No participants met criteria for either Antisocial or Borderline Personality Disorder, and 16 participants from our final sample ($N = 80$) met criteria for at least one Axis I disorder. Given that the exclusion of these individuals did not substantially alter our results, we present data from the entire sample in the main text (specific diagnoses available upon request). In addition, all analyses were conducted both with and without current diagnosis as a covariate (dummy coded: 0 = no psychopathology, 1 = meeting criteria for one or more psychiatric disorders).

Replication cohort. Our replication cohort was drawn from among children and adolescents ($N = 323$, 11–15 years old) participating in the Teen Alcohol Outcomes Study (TAOS) at the University of Texas Health Science Center at San Antonio (UTHSCSA). This ongoing longitudinal study aims to investigate how individual differences in genetic background, environmental experience, and neural function contribute to the emergence of psychopathology, with an emphasis on alcohol use disorders. The current analysis focused on 96 participants (48 girls, mean age 13.62 ± 0.99) who were of Caucasian origin and who had high-quality fMRI data in the first wave of neuroimaging. The study was approved by the institutional review board at UTHSCSA. Consent for study participation was obtained from participants' parents or guardians. Under-age participants provided assent.

Postmortem cohort. The demographic and clinical parameters of the cohort and technical parameters of the samples have been described in detail previously²⁰. All procedures involving this cohort were approved by the University of Pittsburgh's Committee for the Oversight of Research Involving the Dead and Institutional Review Board for Biomedical Research. Consent was obtained from each subject's next of kin. For all subjects, consensus DSM-IV diagnoses of MDD were made by an independent committee of experienced clinical research scientists at a case conference using information obtained from clinical records, toxicology exam and a standardized psychological autopsy²¹. The latter incorporates a structured interview, conducted by a licensed clinical psychologist with family members of the index subject, to assess diagnosis, psychopathology, medical, social and family histories, as well as history of substance abuse. All subjects died suddenly, without prolonged agonal periods. For consistency across samples and to minimize population stratification and pharmacologic confounds, we focused our analysis on Caucasian individuals and excluded those with ascertained medication use at time of death (ascertained by toxicology screen on peripheral tissue), leaving a final sample of 35 (10 women, mean age = 49.57 ± 11.87 years, range = 22–69 years). Brains were analyzed for adequate pH (>6.0) and RNA integrity by optical density (RNA ratio, $OD \geq 1.3$) and Agilent bioanalyzer analysis (Agilent Technologies; RIN expert scoring system ≥ 7) as described²⁰. Gender, age, MDD versus control status, pH, RNA ratio, and post mortem interval were controlled for in all analyses involving this sample. A square root transformation was applied

to *SLC6A4* mRNA levels to normalize its positively skewed and kurtotic distribution (pre-transformation skewness = 1.92, kurtosis = 6.45, post-transformation skewness = 0.65, kurtosis = 1.43).

BOLD fMRI task. As described previously²², the amygdala reactivity procedure used in the discovery cohort (DNS) consisted of four blocks of a face-processing task interleaved with five blocks of a sensorimotor control task. During task blocks, participants viewed a trio of faces (with neutral, angry, fearful or surprised expressions) and matched 1 of 2 faces (bottom) identical to a target face (top). During control blocks, participants matched simple geometric shapes. In the replication cohort (TAOS), the task consisted only of angry and fearful faces. Thus, for consistency between samples, we focused our analyses on the anger + fear > shapes contrast in our discovery cohort. Performance was monitored and participants with accuracy < 75% were excluded from analysis.

BOLD fMRI data acquisition. Participants in the discovery cohort were scanned using a research-dedicated GE MR750 3T scanner equipped with high-power high-duty cycle 50-mT m^{-1} gradients at $200\text{ T m}^{-1}\text{ s}^{-1}$ slew rate, and an eight-channel head coil for parallel imaging at high bandwidth up to 1 MHz at the Duke-UNC Brain Imaging and Analysis Center. A semi-automated high-order shimming program was used to ensure global field homogeneity. A series of 34 interleaved axial functional slices aligned with the anterior commissure-posterior commissure (AC-PC) plane were acquired for full-brain coverage using an inverse-spiral pulse sequence to reduce susceptibility artifact (TR = 2,000 ms; TE = 30 ms; flip angle = 60; FOV = 240 mm; $3.75 \times 3.75 \times 4$ mm voxels; interslice skip = 0). Four initial RF excitations were performed (and discarded) to achieve steady-state equilibrium. To allow for spatial registration of each participant's data to a standard coordinate system, high-resolution three-dimensional structural images were acquired in 34 axial slices co-planar with the functional scans (TR = 7,700 ms; TE = 3.0 ms; flip angle = 12; voxel size = $0.9 \times 0.9 \times 4$ mm; FOV = 240 mm; interslice skip = 0).

Participants in the replication cohort were scanned on a Siemens 3T Trio Scanner at the UTHSCSA. BOLD fMRI data were acquired with a gradient-echo echo planar imaging (EPI) sequence (TR = 2,000 ms; TE = 25 ms; flip angle = 70; FOV = 256 mm; $2.00 \times 2.00 \times 3.00$ mm voxels; interslice skip = 0) covering 34 interleaved 3-mm-thick axial slices. As in the discovery cohort, high-resolution three-dimensional structural images were acquired in 34 axial slices co-planar with the functional scans (TR = 5,610 ms; TE = 72 ms; flip angle = 150; voxel size = $0.8 \times 0.8 \times 3$ mm; FOV = 220 mm \times 320 mm; interslice skip = 0).

BOLD fMRI data preprocessing. The same data preprocessing steps were applied to both the discovery and the replication cohort. Briefly, images for each subject were realigned to the first volume in the time series to correct for head motion, spatially normalized into a standard stereotaxic space (Montreal Neurological Institute template) using a 12-parameter affine model (final resolution of functional images = 2 mm isotropic voxels), and smoothed to minimize noise and residual difference in gyral anatomy with a Gaussian filter, set at 6-mm full-width at half-maximum. Voxel-wise signal intensities were ratio normalized to the whole-brain global mean.

Variability in single-subject whole-brain functional volumes was determined using the Artifact Recognition Toolbox (http://www.nitrc.org/projects/artifact_detect). Individual whole-brain BOLD fMRI volumes meeting at least one of two criteria were assigned a lower weight in determination of task-specific effects: significant mean-volume signal intensity variation (that is, within volume mean signal greater or less than 4 s.d. of mean signal of all volumes in time series), and individual volumes where scan-to-scan movement exceeded 2 mm translation or 2° rotation in any direction.

BOLD fMRI data analysis. The general linear model (GLM) of SPM8 (<http://www.fil.ion.ucl.ac.uk/spm>) was used to conduct fMRI data analyses. Linear contrasts employing canonical hemodynamic response functions were used to estimate differential effects of condition from the contrast of faces > shapes for each individual. Individual contrast images were then used in second-level random effects models accounting for scan-to-scan and participant-to-participant variability to determine mean condition-specific regional responses using one-sample *t* tests. Regions of interest masks for the bilateral amygdala were constructed using the automatic anatomical labeling (AAL) in WFU PickAtlas

Tool, version 1.04. A statistical threshold of $P < 0.05$, FWE corrected, and ≥ 10 contiguous voxels was applied to amygdala analyses within each hemisphere. BOLD values from voxels within the amygdala exhibiting strongest main effect of task were extracted using the VOI tool in SPM8.

These extracted values were then entered into regression models using IBM SPSS Statistics 20.0 (SPSS). Notably, by extracting amygdala BOLD parameter estimates from the voxels activated by our procedure rather than clusters specifically correlated with our independent variables of interest, we preclude the possibility of any regression coefficient inflation that may result from capitalizing on the same data twice²³. We have successfully used this conservative strategy in previous reports^{24–26}.

RNA processing and quantitative real-time PCR. Total RNA was isolated from TRIzol homogenates of the amygdala in all postmortem subjects. The samples were purified using RNeasy spin columns (Qiagen), and RNA integrity was assessed using the Agilent 2100 Bioanalyzer (Agilent Technologies). cDNA was generated by mixing 1 μ g total RNA with oligo-dT primers and SuperScript II reverse transcriptase (Invitrogen) per the manufacturer's protocol. PCR products were amplified in quadruplets on a Mastercycler real-time PCR machine (Eppendorf) using universal PCR conditions, as described previously²⁷. Results were calculated as the geometric mean of threshold cycles of *SLC6A4* transcript amplification normalized to three validated internal controls (actin, glyceraldehyde-3-phosphate dehydrogenase and cyclophilin G). Although performed in serotonergic projection areas, *SLC6A4* transcripts are readily detectable by quantitative PCR.

DNA extraction and 5-HTTLPR/rs25531 genotyping. Saliva samples from discovery cohort participants were collected using Oragene kits and DNA was extracted in accordance with the manufacturer's guidelines (Oragene, Genotek). In the replication cohort, DNA was extracted from whole blood. Postmortem brain DNA was isolated using the DNeasy Blood and Tissue kit (Qiagen), using a protocol that was modified from manufacturers' instructions (additional proteinase K and RNase A).

The same 5-HTTLPR/rs25531 genotyping protocol was applied to DNA samples from all three cohorts. Primer sequences for genotyping 5-HTTLPR were described previously²⁸ (forward: 5'-ATGCCAGCACCTAACCCTAATGT-3'; reverse: 5'-GGACCGCAAGGTGGGCGGGA-3'). PCR was conducted using the following cycling conditions: initial 15-min denaturing step at 95 °C, followed by 35 cycles of 94 °C for 30 s, 66 °C for 30 s and 72 °C for 40 s, and a final extension phase of 72 °C for 15 min. Reactions were performed in 10 \times reaction Buffer IV (ABgene), 1.5 mM MgCl₂, 50 ng of genomic DNA, 5 pmol of each primer, 0.3 mM dNTPs and 1 unit of Native Taq (Promega). PCR products were subsequently digested by MspI restriction enzyme for 4 h at 37 °C. The digestion products were separated on a 3% agarose gel (MultiABgarose, ABgene) supplemented with ethidium bromide (0.03%, vol/vol, BDH) and visualized by ultraviolet trans-illumination. Genotype calls were made by three independent raters, who reached consensus on 100% of the discovery and replication cohort samples. Genotype could not be determined accurately for one post-mortem sample. Thus, it was removed from analysis, leaving a final sample of 34 individuals (10 women, mean age = 49.44 \pm 12.02 years).

DNA methylation analyses by quantitative bisulfite pyrosequencing. DNA methylation levels of the proximal promoter of the serotonin transporter gene in the discovery cohort were determined using quantitative bisulfite pyrosequencing by EpigenDx. Briefly, the human serotonin transport gene (*SLC6A4*) proximal promoter methylation assays analyze 20 CpG dinucleotides in the promoter region from -213 to -69 base pairs of the TSS, based on Ensembl Gene ID [ENSG00000108576](#) and the Transcript ID [ENST00000394821](#). The *SLC6A4* promoter assays (ADS580-FS1 and ADS580-FS2) are targeted to the antisense sequence of *SLC6A4* gene. The targeted CpG loci are listed in **Supplementary Table 1**, along with their genomic location and position relative to the translational start site. The target sequences (genomic DNA and bisulfite converted DNA) are available upon request.

For each analysis, the bisulfite conversion was performed with 500 ng provided genomic DNA using the EZ DNA methylation kit (ZymoResearch). The PCR reaction was performed based on recommended assay conditions (EpigenDx) using 0.2 μ M of each primer with one of the PCR primers being

biotinylated in order to purify the final PCR product using Sepharose beads. The PCR product was bound to Streptavidin Sepharose HP (Amersham Biosciences), and the Sepharose beads containing the immobilized PCR product were purified, washed and denatured using 0.2 M NaOH solution and rewashed using the Pyrosequencing Vacuum Prep Tool (Pyrosequencing, Qiagen) as recommended by the manufacturer. 10 μ l of the PCR products were sequenced by Pyrosequencing PSQ96 HS System (Pyrosequencing, Qiagen) following the manufacturer's instructions (Pyrosequencing, Qiagen). The methylation status of each CpG site was analyzed individually as an artificial T/C single-nucleotide polymorphism using QCpG software (Pyrosequencing, Qiagen).

The methylation level at each CpG site was calculated as the percentage of the methylated alleles over the sum of methylated and unmethylated alleles. The mean methylation level was calculated using methylation levels of all measured CpG sites in each targeted region. For quality control, each experiment included non-CpG cytosines as internal controls to verify efficient sodium bisulfite DNA conversion. We also included low, medium and high methylated standards (EpigenDx) as controls in each run. In light of the low methylation values observed at some CpG sites (<4%) in the discovery cohort, additional PCR bias testing was performed by EpigenDx for the entire ADS580 assay (that is, all 20 CpG sites taken together), as well as individually for CpG 14. The bias testing was conducted by mixing the unmethylated DNA control and *in vitro* methylated DNA at different ratios (0%, 2.5%, 5%, 7.5%, 10%, 50% and 100%) followed by bisulfite modification, PCR and pyrosequencing analysis, which were run in triplicate. There was a high correlation between the percent methylation obtained from the mixing study and expected methylation percentages ($r^2 > 0.97$), which confirms the quality of our data (**Supplementary Fig. 1**).

In the replication and postmortem cohorts, methylation analysis on the same *SLC6A4* proximal promoter 20 CpG sites as targeted by the ADS580FS1 and ADS580FS2 assays was carried out at the Core for Advanced Translational Technologies (UTHSCSA). The protocol used was the same as in the discovery cohort, except with independently designed PCR and sequencing primers (primer sequences available upon request). Results were analyzed using PyroMark Q96 MD and PyroMark CpG 1.0 software (Qiagen).

Control analyses conducted by EpigenDx only in the discovery cohort used additional methylation assays in exon 1 and intron 1 of *SLC6A4* (**Supplementary Table 1**), as well as the promoter region of the *COMT* gene (**Supplementary Table 4**). Detailed assay information is available upon request.

Self-report measures. To assess recent life stress, we administered a modified version of the Life Events Scale for Students (LESS²⁹). This modified version of the scale asks participants to indicate whether they experienced common stressful life events in the past 12 months; in addition, for each event that occurred, participants reported on the impact it had on their lives on a 1–4 scale (with 4 being the highest). The impact scores were set to zero for events that did not occur. Based on prior research^{24,29}, we focused on the LESS Highest Impact metric, reflecting the highest impact associated with any event which occurred within the past year. We assessed early life trauma using the Childhood Trauma Questionnaire (CTQ³⁰).

Statistical analysis. Percent methylation was computed as the ratio of methylation cytosines over the sum of all methylation and unmethylated cytosines. Our main analyses focused on two *SLC6A4* promoter assays covering a total of 20 CpG sites sampled across all three cohorts (**Table 1** and **Supplementary Table 1**).

No single-nucleotide polymorphisms resulting in CpG site gain or loss were identified in the assayed regions. In addition to analysis using average percent methylation across the 20 CpG sites in our region of interest, we applied principal component analysis (PCA) to these 20 CpG sites in both the discovery and replication cohorts. The unrotated correlation matrix was analyzed to output principal component scores. An eigenvalue greater than 1 indicates that PCs account for more variance than accounted for by one of the original variables in standardized data. PCA resulted in five PCs with eigenvalues > 1 in both samples. The first PC exhibited effects similar to those associated with *SLC6A4* promoter methylation values averaged across all 20 CpG sites in both the discovery and the replication cohort (**Supplementary Tables 2, 3, 5 and 6**, and **Supplementary Fig. 1**). In light of the smaller number of data points ($n = 34$ final sample), no PCA was performed in the postmortem cohort, where CpG sites were analyzed individually.

Linear regression models, as implemented in IBM SPSS Statistics 20.0, were used to investigate the linear effect of methylation values (independent variable) on amygdala reactivity or *SLC6A4* mRNA levels (dependent variables). Results from two-tailed tests are reported for all analyses.

A **Supplementary Methods Checklist** is available.

18. Sheehan, D.V. *et al. J. Clin. Psychiatry* **59** (suppl. 20), 22–33, quiz 34–57 (1998).
19. First, M.B., Spitzer, R.L., Gibbon, M. & Williams, J.B.M. *Structured Clinical Interview for DSM-IV Axis I Disorders, Research Version, Non-patient Edition* (New York State Psychiatric Institute, Biometrics Research Department, New York, 1996).
20. Sibille, E. *et al. Am. J. Psychiatry* **166**, 1011–1024 (2009).
21. Glantz, L.A. & Lewis, D.A. *Arch. Gen. Psychiatry* **54**, 943–952 (1997).
22. Carré, J.M., Hyde, L.W., Neumann, C.S., Viding, E. & Hariri, A.R. *Soc. Neurosci.* **8**, 122–135 (2013).
23. Viviani, R. *Neuroimage* **50**, 184–189 (2010).
24. Nikolova, Y.S. & Hariri, A.R. *Biol. Mood Anxiety Disord.* **2**, 19 (2012).
25. Nikolova, Y.S., Ferrell, R.E., Manuck, S.B. & Hariri, A.R. *Neuropsychopharmacology* **36**, 1940–1947 (2011).
26. Carré, J.M., Fisher, P.M., Manuck, S.B. & Hariri, A.R. *Soc. Cogn. Affect. Neurosci.* **7**, 213–221 (2012).
27. Tripp, A. *et al. Am. J. Psychiatry* **169**, 1194–1202 (2012).
28. Gelernter, J., Kranzler, H. & Cubells, J.F. *Hum. Genet.* **101**, 243–246 (1997).
29. Nikolova, Y.S., Bogdan, R., Brigidi, B.D. & Hariri, A.R. *Biol. Psychiatry* **72**, 157–163 (2012).
30. Bernstein, D.P. *et al. Child Abuse Negl.* **27**, 169–190 (2003).